

To help adults control type 2 diabetes along with diet and exercise

# ALONG WITH DIET AND EXERCISE, CYCLOSET MAY LOWER BLOOD SUGAR LEVELS IN ADULTS WITH TYPE 2 DIABETES.

Taking CYCLOSET as part of your  
type 2 diabetes treatment regimen

**CO-PAY SAVINGS INFORMATION INSIDE!**

## **INDICATION**

CYCLOSET® (bromocriptine mesylate) 0.8 mg tablets is a once-daily prescription pill used with diet and exercise to lower blood sugar levels in adults with type 2 diabetes.

CYCLOSET is not for people with type 1 diabetes or for people with diabetic ketoacidosis. It is not known if CYCLOSET is safe and effective in children.

## **IMPORTANT SAFETY INFORMATION**

### **Do not take CYCLOSET if you:**

- Are allergic to bromocriptine or any of the other ingredients in CYCLOSET.
- Are allergic to ergot medicines. Ask your health care provider or pharmacist for a list of these medicines, if you are not sure whether you take any.

**Please see additional  
Important Safety Information  
throughout and accompanying  
full Prescribing Information.**

  
**CYCLOSET**®  
bromocriptine mesylate tablets

## CYCLOSET LOWERS BLOOD SUGAR

CYCLOSET is an oral once-daily prescription medicine that helps lower blood sugar levels in adults with type 2 diabetes, when used with a healthy diet and exercise. It may help you reach your blood sugar goals you set with your doctor.

There are two major problems causing type 2 diabetes:

- The body does not produce enough insulin to adequately control blood glucose levels
- The body becomes insulin resistant, meaning it cannot use insulin effectively

Your doctor has prescribed CYCLOSET as an important part of your care to help your body use insulin and lower your blood sugar after meals.

## HOW DOES CYCLOSET WORK?

- CYCLOSET is a prescription medicine used with diet and exercise to lower blood sugar in adults with type 2 diabetes
- In clinical studies, CYCLOSET therapy has been shown to lower blood sugar levels after each meal of the day and improve HbA1c levels
- CYCLOSET, taken once a day, in the morning, within 2 hours of waking up, with food, lowered blood sugar levels at 2 hours after each meal, without raising insulin levels in the blood

## IMPORTANT SAFETY INFORMATION (cont'd)

### Do not take CYCLOSET if you:

- Have fainting migraine headaches.
- Are a woman who has recently given birth to a child (postpartum). CYCLOSET can cause serious and life-threatening problems including high blood pressure (hypertension), heart attack, seizures, stroke, and psychosis (losing touch with reality).
- Are breastfeeding.





### CYCLOSET may cause serious side effects, including:

- **Low blood pressure, fainting, or severe dizziness after you stand up from a lying down position.** Stand up slowly when moving from a sitting or lying position. You should avoid activities that could cause injury if you feel light-headed or faint.
- **Feeling tired and sleepy.** If you feel tired and sleepy while taking CYCLOSET, you should not drive or operate heavy machinery.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information.







## WHAT TO REMEMBER WHEN TAKING CYCLOSET


### CYCLOSET should be taken once daily:

-  In the morning, within 2 hours of waking up
-  With food (your morning meal)
-  If you miss a dose, just take your usual dose the next morning
-  **DO NOT** take a double dose of CYCLOSET

Your doctor will work on a dosing schedule with you and tell you how many CYCLOSET tablets to take each day.

The following page has a handy chart that you can use to help keep track of your CYCLOSET dose.

RECOMMENDED DOSING SCHEDULE						
WEEK	1	2	3	4	5	6
# OF TABLETS PER DAY	1 	2 	3 	4 	5 	6 

 Actual tablet size

- Your doctor may increase your daily morning dose by 1 tablet each week, **until you reach an appropriate dose**, typically between 2 and 6 tablets per day
  - Your doctor may tell you to increase your dose more slowly

While you are taking CYCLOSET, remember to:

- Stay on your prescribed diet and exercise program
- Check your blood sugar 1 to 2 hours after each meal or as directed by your doctor
- Share your tracking sheet with your doctor at your next appointment to review your treatment progress

# SET UP YOUR CYCLOSET® DOSING SCHEDULE

## SAVE ON CYCLOSET

### PARTNER WITH YOUR DOCTOR ON YOUR DOSING PLAN

Use this medication tracker to make sure you take your dose of CYCLOSET as prescribed each day.

WEEK	# OF TABLETS PER DAY	TIME CYCLOSET TAKEN						
		DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
1								
2								
3								
4								
5								
6								
7								
8								
9								

### IMPORTANT SAFETY INFORMATION (cont'd)

#### CYCLOSET may cause serious side effects, including:

- Uncontrollable (compulsive) behaviors that include strong urges to gamble, increased sexual urges, strong urges to spend money, binge eating or eating that you cannot control. These behaviors may go away if your dose of CYCLOSET is lowered or stopped.
- **Serious and life-threatening problems in women who recently have given birth to a child.**

#### Before taking CYCLOSET, tell your health care provider about all medical conditions, including if you:

- Have type 1 diabetes mellitus or diabetic ketoacidosis.
- Have ever passed out or fainted.
- Have migraine headaches.
- Have or have had low blood pressure (hypotension).
- Take medicines for high blood pressure (hypertension).
- Have or have had a mental health condition, especially a psychotic disorder.
- Are pregnant or plan to become pregnant. It is not known if CYCLOSET will harm your unborn baby. Talk with your health care provider if you are pregnant or plan to become pregnant.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

### THE CYCLOSET CO-PAY SAVINGS PROGRAM IS BUILT WITH PATIENTS IN MIND

With the CYCLOSET Co-Pay Savings Program, you may qualify to pay as little as a \$0\* co-pay per prescription fill (maximum benefits apply\*\*).

For more information, call **1-866-686-0049** or visit [CYCLOSET.com](#).

With the CYCLOSET Co-Pay Savings Program  
**MOST COMMERCIALY INSURED PATIENTS MAY PAY AS LITTLE AS A**

**\$0\*** **CO-PAY**  
1<sup>ST</sup> Rx AND REFILLS

DOWNLOAD THE COUPON AT [CYCLOSET.COM](#)

\*This offer is only valid for patients with commercial insurance. Uninsured patients are not eligible. This offer is not valid for any person eligible for reimbursement of prescriptions, in whole or in part, by any federal, state, or other governmental programs, including, but not limited to, Medicare (including Medicare Advantage and Part A, B, and D plans), Medicaid, TRICARE, Veterans Administration or Department of Defense health coverage, CHAMPUS, the Puerto Rico Government Health Insurance Plan, or any other federal or state health care programs. This offer is not valid in Massachusetts or Minnesota or where otherwise prohibited, taxed or otherwise restricted. Please see [CYCLOSET.com](#) for full Eligibility Terms and Conditions.

\*\*Patient is responsible for all additional costs and expenses after application of the maximum benefits. For information about the maximum benefits, please call 1-866-686-0049.

### IMPORTANT SAFETY INFORMATION (cont'd)

#### Tell your health care provider about all medicines you take and any recent changes to your medications (prescription, non-prescription, vitamins and herbal supplements), especially those for:

- Mental health conditions, especially anti-psychotic medicines called dopamine receptor antagonists.
- Migraine or other types of headaches.
- Other medicines that work like CYCLOSET (dopamine receptor agonists) to treat conditions such as Parkinson's disease, high levels of a hormone called prolactin in your blood (hyperprolactinemia), restless leg syndrome and too much growth hormone (acromegaly).
- Type 2 diabetes.



## STAY ON TRACK

**It is important to track your treatment progress and share it with your doctor at your appointments.**

### **IMPORTANT SAFETY INFORMATION (cont'd)**

#### **The most common side effects of CYCLOSET were:**

- Nausea, headache, dizziness, vomiting, and/or lightheadedness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch/](http://www.fda.gov/medwatch/) or call 1-800-FDA-1088.

For product information, adverse event reports, and product complaint reports, please contact:

Salix Product Information Call Center

Phone: 1-800-321-4576

Fax: 1-510-595-8183

Email: [salixmc@dlss.com](mailto:salixmc@dlss.com)

**Please see additional Important Safety Information throughout and accompanying full Prescribing Information.**



The Salix logo is a trademark of Salix Pharmaceuticals or its affiliates. CYCLOSET is a registered trademark of VeroScience LLC, Tiverton, RI 02878 used under license. ©2021 Salix Pharmaceuticals or its affiliates. CYC.0072.USA.20 V3.0.